

Prevention Strategies

Lisa A. Kisling; Joe M. Das.

[Author Information and Affiliations](#)

Last Update: August 1, 2023.

[:](#)

Definition/Introduction

The natural history of a disease classifies into five stages: underlying, susceptible, subclinical, clinical, and recovery/disability/death. Corresponding preventive health measures have been grouped into similar stages to target the prevention of these stages of a disease. These preventive stages are primordial prevention, primary prevention, secondary prevention, and tertiary prevention. Combined, these strategies not only aim to prevent the onset of disease through risk reduction but also downstream complications of a manifested disease.

Primordial Prevention

In 1978, the most recent addition to preventive strategies, **primordial prevention**, was described. It consists of risk factor reduction targeted towards an entire population through a focus on social and environmental conditions. Such measures typically get promoted through laws and national policy. Because primordial prevention is the earliest prevention modality, it is often aimed at children to decrease as much risk exposure as possible. Primordial prevention targets the underlying stage of natural disease by targeting the underlying social conditions that promote disease onset. An example includes improving access to an urban neighborhood to safe sidewalks to promote physical activity; this, in turn, decreases risk factors for obesity, cardiovascular disease, type 2 diabetes, etc.

Primary Prevention

Primary prevention consists of measures aimed at a susceptible population or individual. The purpose of primary prevention is to prevent a disease from ever occurring. Thus, its target population is healthy individuals. It commonly institutes activities that limit risk exposure or increase the immunity of individuals at risk to prevent a disease from progressing in a susceptible individual to subclinical disease. For example, immunizations are a form of primary prevention.

Secondary Prevention

Secondary prevention emphasizes early disease detection, and its target is healthy-appearing individuals with subclinical forms of the disease. The subclinical disease consists of pathologic changes but no overt symptoms that are diagnosable in a doctor's visit. Secondary prevention often occurs in the form of screenings. For example, a Papanicolaou (Pap) smear is a form of secondary prevention aimed to diagnose cervical cancer in its subclinical state before progression.

Tertiary Prevention

Tertiary prevention targets both the clinical and outcome stages of a disease. It is implemented in symptomatic patients and aims to reduce the severity of the disease as well as any associated sequelae. While secondary prevention seeks to prevent the onset of illness, tertiary prevention aims to reduce the effects of the disease once established in an individual. Forms of tertiary prevention are commonly rehabilitation efforts.

Quaternary Prevention

According to the Wonca International Dictionary for General/Family Practice, quaternary prevention is "action taken to identify patients at risk of overmedicalization, to

protect him from new medical invasion, and to suggest to him interventions, which are ethically acceptable." Marc Jamoulle initially proposed this concept, and the targets were mainly patients with illness but without the disease. The definition has undergone recent modification as "an action taken to protect individuals (persons/patients) from medical interventions that are likely to cause more harm than good." [\[1\]](#)

[Go to:](#)

Issues of Concern

In the United States, several governing bodies make prevention recommendations. For example, the United States Preventive Services Task Force (USPSTF) is a governing body that makes recommendations for primary and secondary prevention strategies. The Advisory Committee on Immunizations Practices (ACIP) through the Centers for Disease Control and Prevention (CDC) makes recommendations for vaccinations, while the Women's Preventive Services Initiative (WPSI) makes recommendations appropriate for females. Additionally, various specialty organizations, such as the American College of Obstetrics and Gynecology (ACOG) and the American Cancer Society (ACS), etc. also make prevention recommendations. With the multitude of information and recommending bodies, it is often challenging for healthcare professionals to remain up to date on changing endorsements.

Further, while preventive services are regulated and must undergo scrupulous safety testing, there is risk involved with prevention. Particularly primary and secondary preventive factors targeted at intervening in healthy-appearing individuals. It is often challenging to gain buy-in from patients regarding the risk-benefit ratio of various preventive services.

Finally, the cost of preventive services is commonly a topic of discussion. Several cost-benefit analyses have been undertaken regarding the evaluation of preventive services with varying degrees of confidence. While often a long-term gain of healthy life years is noted, preventive services are not inexpensive, which can limit the use of these services by both healthcare systems and patients and is a consideration when promoting preventive services.

[Go to:](#)

Clinical Significance

Preventive services have proven an essential aspect of healthcare; however, they appear consistently underutilized in the United States.[\[2\]](#)[\[3\]](#)[\[4\]](#) With cost, time, and resource constraints on physicians, many preventive services get overlooked. Physicians need to remain up to date on the prevention guidelines and ensure all patients are offered appropriate services with a full explanation of risks and benefits.

Some examples of commonly used prevention strategies are:

Primordial

- Government policy: Increasing taxes on cigarettes; Decreasing advertisement of tobacco[\[5\]](#)
- Built Environment: Access to safe walking paths; access to stores with healthy food options

Primary

- Immunizations[\[6\]](#)
- Tobacco cessation programs
- Needle exchange programs[\[7\]](#)
- Micronutrient supplementation programs

Secondary

- Papanicolaou (Pap) smear for early detection of cervical cancer[\[8\]](#)
- Mammography for early detection of breast cancer
- Colonoscopies for early detection of colon cancer
- Blood Pressure Screening

Tertiary

- Occupational and physical therapy in burn patients
- Cardiac rehab in post-myocardial infarction patients
- Diabetic foot care

Quarternary

The following conditions are susceptible to over-treatment:

- Radiological incidentalomas[\[9\]](#)
- The use of antiarrhythmic drugs after myocardial infarction that reduced arrhythmias but increased mortality
- The use of hormone replacement therapy led to an increased number of cases of breast cancer, stroke, and thromboembolic events. It was also a failure in reducing cardiovascular mortality.
- Medically unexplained symptoms
- Functional disorders
- Bodily distress syndrome[\[10\]](#)

[Go to:](#)

Nursing, Allied Health, and Interprofessional Team Interventions

Proper communication among the various healthcare personnel should be there to provide appropriate levels of

prevention to the general public and patients. School staff and other ancillary staff require education on the importance of providing prevention as an important aspect of caring for an individual while he/she is a student.

[Go to:](#)

Review Questions

- [Access free multiple choice questions on this topic.](#)
- [Comment on this article.](#)

[Go to:](#)

References

1. Martins C, Godycki-Cwirko M, Heleno B, Brodersen J. Quaternary prevention: reviewing the concept. Eur J Gen Pract. 2018 Dec;24(1):106-111. [[PMC free article](#)] [[PubMed](#)]
2. Chung S, Romanelli RJ, Stults CD, Luft HS. Preventive visit among older adults with Medicare's introduction of Annual Wellness Visit: Closing gaps in underutilization. Prev Med. 2018 Oct;115:110-118. [[PMC free article](#)] [[PubMed](#)]
3. Kottke TE, Solberg LI, Brekke ML, Cabrera A, Marquez MA. Delivery rates for preventive services in 44 midwestern clinics. Mayo Clin Proc. 1997 Jun;72(6):515-23. [[PubMed](#)]
4. Smith RA, Andrews KS, Brooks D, Fedewa SA, Manassaram-Baptiste D, Saslow D, Brawley OW, Wender RC. Cancer screening in the United States, 2017: A review of current American Cancer Society

guidelines and current issues in cancer screening. CA Cancer J Clin. 2017 Mar;67(2):100-121. [[PubMed](#)]

5.

Fielding R, Chee YY, Choi KM, Chu TK, Kato K, Lam SK, Sin KL, Tang KT, Wong HM, Wong KM. Declines in tobacco brand recognition and ever-smoking rates among young children following restrictions on tobacco advertisements in Hong Kong. J Public Health (Oxf). 2004 Mar;26(1):24-30. [[PubMed](#)]

6.

Maier C, Maier T, Neagu CE, Vlădăreanu R. Romanian adolescents' knowledge and attitudes towards human papillomavirus infection and prophylactic vaccination. Eur J Obstet Gynecol Reprod Biol. 2015 Dec;195:77-82. [[PubMed](#)]

7.

Calvo F, Carbonell X, Rived M, Giralt C. When people who inject drugs speak: Qualitative thematic analysis of the perception of a mobile app for needle exchange programs. Adicciones. 2021 Jul 01;33(3):217-234. [[PubMed](#)]

8.

Nguyen TT, McPhee SJ, Nguyen T, Lam T, Mock J. Predictors of cervical Pap smear screening awareness, intention, and receipt among Vietnamese-American women. Am J Prev Med. 2002 Oct;23(3):207-14. [[PMC free article](#)] [[PubMed](#)]

9.

Sherlock M, Scarsbrook A, Abbas A, Fraser S, Limumpornpetch P, Dineen R, Stewart PM. Adrenal Incidentaloma. Endocr Rev. 2020 Dec 01;41(6):775-820. [[PMC free article](#)] [[PubMed](#)]

10.

Schmalbach B, Roenneberg C, Hausteiner-Wiehle C, Henningsen P, Brähler E, Zenger M, Häuser W.

Validation of the German version of the Bodily Distress Syndrome 25 checklist in a representative German population sample. J Psychosom Res. 2020 May;132:109991. [[PubMed](#)]

Disclosure: Lisa Kisling declares no relevant financial relationships with ineligible companies.

Disclosure: Joe Das declares no relevant financial relationships with ineligible companies.

[Copyright](#) © 2024, StatPearls Publishing LLC.

This book is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits others to distribute the work, provided that the article is not altered or used commercially. You are not required to obtain permission to distribute this article, provided that you credit the author and journal.

Bookshelf ID: NBK537222PMID: [30725907](#)

- [Share on Facebook](#)
- [Share on Twitter](#)

Views

- [PubReader](#)
 - [Print View](#)
 - [Cite this Page](#)
-

In this Page

- [Definition/Introduction](#)
- [Issues of Concern](#)
- [Clinical Significance](#)

- [Nursing, Allied Health, and Interprofessional Team Interventions](#)
 - [Review Questions](#)
 - [References](#)
-

Bulk Download

- [Bulk download StatPearls data from FTP](#)
-

Related information

- [PMC](#)
 - [PubMed](#)
-

Similar articles in PubMed

- [Review Unveiling the Significance and Challenges of Integrating Prevention Levels in Healthcare Practice.](#)[J Prim Care Community Health. ...]
 - [Review Skin cancer: Primary, secondary, and tertiary prevention. Part I.](#)[J Am Acad Dermatol. 2022]
 - [Early prevention of childhood disability in developing countries.](#)[Int J Rehabil Res. 1991]
 - [Review Primary prevention of coronary heart disease: integration of new data, evolving views, revised goals, and role of rosuvastatin in management. A comprehensive survey.](#)[Drug Des Devel Ther. 2011]
 - [Review Prevention of stroke: a global perspective.](#)[Lancet. 2018]
-

[See reviews...](#)[See all...](#)

Recent Activity

[ClearTurn Off](#)

- [Prevention Strategies - StatPearls](#)
 - [Polymerase Chain Reaction \(PCR\) - StatPearls](#)
-

[See more...](#)

FO